

## Continuity and Compactness

Def. A mapping  $f: E \subseteq X \rightarrow \mathbb{R}^k$  is said to be **bounded** if there exists a real number  $M$  such that  $\|f(x)\| \leq M$  for all  $x \in E$ .

Ex.  $f(x, y) = x^2 + y^2$  is bounded for  $E = \{(x, y) \mid |x| < 10, |y| \leq 5\}$

since  $|f(x, y)| \leq 100 + 25 = 125 = M$ ;


But it is not bounded on  $E = \mathbb{R}^2$ .

Ex.  $f(x, y) = e^{-(x^2+y^2)}$  is bounded for  $E = \mathbb{R}^2$  since

$$|f(x, y)| = |e^{-(x^2+y^2)}| \leq 1 = M.$$

Theorem: Suppose  $f: X \rightarrow Y$  is a continuous mapping of a compact metric space  $X$  into a metric space  $Y$  then  $f(X)$  is compact.

Proof: Let  $\{V_\alpha\}$  be an open cover of  $f(X)$ .



Since  $f$  is continuous,  $f^{-1}(V_\alpha)$ , is an open set in  $X$  (why?) and  $X \subseteq \bigcup_\alpha f^{-1}(V_\alpha)$ .

Thus  $\{f^{-1}(V_\alpha)\}$  is an open cover of  $X$ .

Since  $X$  is compact there exists a finite subcover:  $X \subseteq \bigcup_{i=1}^n f^{-1}(W_i)$ , where

$$\{W_i\} \subseteq \{V_\alpha\}.$$

Since  $f(f^{-1}(E)) \subseteq E$ ; for  $E \subseteq Y$ ,

(For example, if  $f(x) = x^2$  and  $E = (-1,1)$ ; then  $f^{-1}(-1,1) = (-1,1)$  and  $f(f^{-1}(-1,1)) = [0,1) \subseteq (-1,1)$ .)

$$f(X) \subseteq \bigcup_{i=1}^n f(f^{-1}(W_i)) \subseteq \bigcup_{i=1}^n W_i .$$

So  $\{W_i\}$  is a finite subcover of  $f(X)$ , and  $f(X)$  is compact.

Theorem: Suppose  $f$  is a continuous function on a compact metric space  $X$  into  $\mathbb{R}$ , and  $M = \sup_{p \in X} f(p)$  and  $m = \inf_{p \in X} f(p)$ , then there exist points  $p, q \in X$  such that  $f(p) = M$  and  $f(q) = m$ .

Proof: Since  $f$  is continuous and  $X$  is compact,  $f(X)$  is a compact subset of  $\mathbb{R}$ . By the Heine-Borel theorem we know that any compact subset of  $\mathbb{R}$  (or  $\mathbb{R}^n$ ) is closed and bounded.

Hence  $f(X)$  contains  $M = \sup_{p \in X} f(p)$  and  $m = \inf_{p \in X} f(p)$ .

For suppose  $M = \sup_{p \in X} f(p)$  and  $M \notin f(X)$ .

Then for every  $h > 0$  there exists a point  $x \in f(X)$  such that  $M - h < x < M$ .

Otherwise,  $M - h$  would be an upper bound for  $f(X)$  and  $M$  wouldn't be the least upper bound.

But this means that  $M$  is a limit point of  $f(X)$ . Since  $f(X)$  is closed  $M \in f(X)$ .

A similar argument works for the infimum.

This gives us the theorem from first year Calculus that a continuous function on a closed, bounded interval (i.e. a compact subset of  $\mathbb{R}$ ) takes on its maximum and minimum values.




Ex. Let  $f(x) = x^2$ ; and  $X = [-1, 1]$ .

$f(X) = [0, 1]$ ; minimum value=0, maximum value=1. (In red below)

If  $X = (-1, 1)$ ; i.e.  $X$  is not compact notice that:


$f(X) = [0, 1)$ ;  $f$  takes on its minimum value but not its maximum value. (In blue below)

If  $X = (0, 1)$ , then  $f(X) = (0, 1)$  and  $f$  doesn't take on either its minimum or maximum values. (In green below)



Note: A continuous function on a non-compact set can take on its minimum and/or maximum values, but it does not have to. A continuous function on a compact set must take on its maximum and minimum values.

Ex. Let  $f(x) = \sin x$ ; and  $X = (0, 2\pi)$ . Then  $f(X) = [-1, 1]$ . So  $f$  takes on its maximum and minimum values even though  $X = (0, 2\pi)$  is not compact.



Def. Let  $f: X \rightarrow Y$ ;  $X, Y$  are metric spaces. We say  $f$  is **uniformly continuous** on  $X$  if for every  $\epsilon > 0$  there exists a  $\delta > 0$  for all  $p, q \in X$  such that if  $d_X(p, q) < \delta$  then  $d_Y(f(p), f(q)) < \epsilon$ .

For any interval  $I \subseteq \mathbb{R}$  with  $f: I \subseteq \mathbb{R} \rightarrow \mathbb{R}$ ,  $f$  is uniformly continuous on  $I$  means for every  $\epsilon > 0$  there exists a  $\delta > 0$  for all  $x, a \in I$  such that if  $|x - a| < \delta$  then  $|f(x) - f(a)| < \epsilon$ .

Notice the difference between continuity and uniform continuity:

1. For uniform continuity,  $\delta$  does not depend on the point in  $X$  you are at. For continuity, the  $\delta$  can depend on which point in  $X$  you are at (with both continuity and uniform continuity,  $\delta$  does depend on  $\epsilon$ ).
2. Uniform continuity is a property of a set of points, not a single point. Continuity is a property at a point and a set of points.
3. If a function is uniformly continuous on a set  $X$ , then it is also continuous on  $X$ . However, if a function is continuous on a set  $X$  it may, or may not be, uniformly continuous on  $X$ .

Ex. Let  $f(x) = \frac{1}{x}$ ;  $0 < x < 1$ . Show that  $f(x)$  is continuous on  $(0,1)$  but not uniformly continuous.

To show  $f(x) = \frac{1}{x}$  is continuous at any point  $a \in (0,1)$  we must show that given any  $\epsilon > 0$  we can find a  $\delta > 0$  such that if  $|x - a| < \delta$  then  $\left| \frac{1}{x} - \frac{1}{a} \right| < \epsilon$ .

Note that  $\delta$  can depend on both the value of  $\epsilon$  and the value of " $a$ ".

Let's work backward from the  $\epsilon$  statement to get the  $\delta$  statement.

$$\left| \frac{1}{x} - \frac{1}{a} \right| = \left| \frac{a-x}{ax} \right| = \frac{1}{|ax|} |x - a|.$$

We need an upper bound on  $\frac{1}{|ax|} = \frac{1}{ax}$ ; since  $a, x > 0$ .

Choose  $\delta \leq \frac{a}{2}$ . (Since  $0 < a < 1$ , we have to choose a  $\delta$  neighborhood that stays away from  $x = 0$ , otherwise  $\frac{1}{ax}$  won't be bounded above).

Then we have that  $|x - a| < \frac{a}{2}$  or

$$-\frac{a}{2} < x - a < \frac{a}{2} \quad \text{now add } a$$

$$\frac{a}{2} < x < \frac{3a}{2};$$

Since  $\frac{a}{2}, x, \frac{3a}{2} > 0$ :  $\frac{2}{a} > \frac{1}{x} > \frac{2}{3a}$ ; now multiply through by  $\frac{1}{a} > 0$ .

$$\frac{2}{a^2} > \frac{1}{ax} > \frac{2}{3a^2} \implies \frac{1}{|ax|} < \frac{2}{a^2}.$$

Thus we have: if  $\delta \leq \frac{a}{2}$  then

$$\left| \frac{1}{x} - \frac{1}{a} \right| = \frac{1}{|ax|} |x - a| < \frac{2}{a^2} |x - a| < \frac{2}{a^2} \delta < \epsilon.$$

$$\frac{2}{a^2} \delta < \epsilon \text{ is equivalent to } \delta < \frac{a^2}{2} \epsilon.$$

Choose  $\delta = \min\left(\frac{a}{2}, \frac{a^2}{2} \epsilon\right)$  (remember we chose  $\delta \leq \frac{a}{2}$  earlier)

Now let's show that this  $\delta$  works.

If  $|x - a| < \delta = \min\left(\frac{a}{2}, \frac{a^2}{2} \epsilon\right)$  then we have:

$$\left| \frac{1}{x} - \frac{1}{a} \right| = \frac{1}{|ax|} |x - a| < \frac{2}{a^2} |x - a| \quad \text{since } \delta \leq \frac{a}{2}$$

$$\left| \frac{1}{x} - \frac{1}{a} \right| < \frac{2}{a^2} |x - a| < \frac{2}{a^2} \delta \leq \frac{2}{a^2} \left(\frac{a^2}{2} \epsilon\right) = \epsilon \quad \text{since } \delta \leq \frac{a^2}{2} \epsilon.$$

Thus we have shown that  $f(x) = \frac{1}{x}$  is continuous at  $a \in (0,1)$ .

Now let's show that  $f(x) = \frac{1}{x}$  is not uniformly continuous on  $(0,1)$ .

Let's fix an  $\epsilon > 0$ .

To be uniformly continuous we need to find a  $\delta > 0$ , that depends only on  $\epsilon$ , such that if  $|x - a| < \delta$  then  $\left| \frac{1}{x} - \frac{1}{a} \right| < \epsilon$  for all  $a, x \in (0,1)$ .

But if  $\epsilon > 0$  is fixed, regardless of what  $\delta$  one chooses, by moving " $a$ " toward 0

$$\left| \frac{1}{x} - \frac{1}{a} \right| = \frac{1}{|ax|} |x - a| \rightarrow \infty \text{ for } |x - a| < \delta.$$

So  $\delta$  must depend on " $a$ " and  $f(x) = \frac{1}{x}$  is not uniformly continuous on  $(0,1)$ .

Ex. Show  $f(x) = x^2$  is uniformly continuous on  $[-1,1]$ .

We must show that given any  $\epsilon > 0$  there exists a  $\delta > 0$  for all  $a, x \in [-1,1]$  such that if  $|x - a| < \delta$  then  $|x^2 - a^2| < \epsilon$ .

Let's start with the  $\epsilon$  statement:

$$|x^2 - a^2| = |x - a||x + a|.$$

But we also know that  $a, x \in [-1,1]$ , so  $|a| \leq 1$  and  $|x| \leq 1$ .

Now using the triangle inequality:  $|x + a| \leq |x| + |a| \leq 1 + 1 = 2$ .

$$\text{So } |x^2 - a^2| = |x - a||x + a| \leq 2|x - a| < 2\delta.$$

So if we can force  $2\delta < \epsilon$ , we'll almost be done.

So if we choose  $\delta < \frac{\epsilon}{2}$  (notice  $\delta$  doesn't depend on  $a$ ) we have:

$$|x^2 - a^2| = |x - a||x + a| \leq 2|x - a| \quad \text{because } |a| \leq 1 \text{ and } |x| \leq 1$$

$$|x^2 - a^2| \leq 2|x - a| < 2\delta < 2\left(\frac{\epsilon}{2}\right) = \epsilon \quad \text{because } \delta < \frac{\epsilon}{2}.$$

Hence  $f(x) = x^2$  is uniformly continuous on  $[-1,1]$ .



Theorem: Let  $f: X \rightarrow Y$ , be continuous,  $X, Y$  metric spaces with  $X$  compact, then  $f$  is uniformly continuous on  $X$ .

Notice that  $f(x) = x^2$  is uniformly continuous on  $(-1,1)$  (as well as on  $[-1,1]$ ) even though  $(-1,1)$  is not compact. The same  $\delta, \epsilon$  argument that shows that  $f(x) = x^2$  is uniformly continuous on  $[-1,1]$  also show that it's uniformly continuous on  $(-1,1)$ . Thus a continuous function on a compact set **must be** uniformly continuous on the compact set. A continuous function on a non-compact set, may or may not be uniformly continuous on that set.

Some special properties of uniformly continuous functions:

1. If  $f: E \rightarrow Y$  is uniformly continuous on a metric space  $E$  and  $\{p_n\}$  is a Cauchy sequence in  $E$ , then  $\{f(p_n)\}$  is a Cauchy sequence in  $Y$ .

Notice that  $f(x) = \frac{1}{x}$  is continuous on  $(0, \infty)$ , but not uniformly continuous.  $\{\frac{1}{n}\}$  is a Cauchy sequence in  $(0, \infty)$ , but  $\{f(\frac{1}{n})\} = \{n\}$  is not.

2. If  $f: E \subseteq \mathbb{R} \rightarrow \mathbb{R}$  is uniformly continuous,  $E$  a bounded interval, then

$\int_E f(x)dx$  is finite.

$f(x) = \frac{1}{x}$  is continuous on  $(0,1)$ , but not uniformly continuous.  $\int_0^1 \frac{1}{x} dx$  is not finite.

Notice also that you can have bounded continuous functions that are not uniformly continuous (e.g.  $f(x) = \sin\left(\frac{1}{x}\right)$ ;  $0 < x < 2\pi$ ) and unbounded continuous functions that are uniformly continuous (e.g.  $f(x) = x$ ;  $-\infty < x < \infty$ ).

Ex. Prove that  $f(x) = \frac{x}{1-x}$  is uniformly continuous on  $[2, \infty)$ .

We must show given any  $\epsilon > 0$  there exists a  $\delta > 0$  for all  $a, x \in [2, \infty)$  such that if  $|x - a| < \delta$  then  $\left| \frac{x}{1-x} - \frac{a}{1-a} \right| < \epsilon$ .

Let's start with the  $\epsilon$  statement:

$$\begin{aligned} \left| \frac{x}{1-x} - \frac{a}{1-a} \right| &= \left| \frac{x(1-a) - a(1-x)}{(1-x)(1-a)} \right| = \left| \frac{(x-a)}{(1-x)(1-a)} \right| \\ &= |x - a| \left| \frac{1}{(1-x)(1-a)} \right|. \end{aligned}$$

Now we must find an upper bound on  $\left| \frac{1}{(1-x)(1-a)} \right|$  independent of "a".

Since  $2 \leq x$  and  $2 \leq a$ :  $-1 \leq \frac{1}{1-x} < 0$  and  $-1 \leq \frac{1}{1-a} < 0$ .

Thus we can say:  $0 < \left(\frac{1}{1-x}\right)\left(\frac{1}{1-a}\right) \leq 1$ .

Thus we have:

$$\left| \frac{x}{1-x} - \frac{a}{1-a} \right| = |x - a| \left| \frac{1}{(1-x)(1-a)} \right| \leq 1|x - a| = \delta.$$

So if we can force  $\delta < \epsilon$  we'll almost be done.

So choose  $\delta < \epsilon$  which is independent of "a".

Now let's show  $\delta < \epsilon$  works.

If  $|x - a| < \delta < \epsilon$  then:

$$\left| \frac{x}{1-x} - \frac{a}{1-a} \right| = |x - a| \left| \frac{1}{(1-x)(1-a)} \right| < |x - a| \quad \text{because } 2 \leq x \text{ and } 2 \leq a$$

$$\left| \frac{x}{1-x} - \frac{a}{1-a} \right| < |x - a| < \delta < \epsilon \quad \text{because } \delta < \epsilon.$$

Hence  $f(x) = \frac{x}{1-x}$  is uniformly continuous on  $[2, \infty)$ .