

Tangent Spaces


Let $M \subseteq \mathbb{R}^n$ be a k -dimensional manifold and $\vec{\Phi}$ a parameterization where $\vec{\Phi}: U \subseteq \mathbb{R}^k \rightarrow M \subseteq \mathbb{R}^n$ and $\vec{\Phi}(a) = x \in M$, then:

$$D\vec{\Phi}(a): \mathbb{R}_a^k \rightarrow \mathbb{R}_x^n.$$

Def. We call $D\vec{\Phi}(a)(\mathbb{R}_a^k) = T_x(M)$ the **tangent space** of M at x .

Note: this definition does not depend on the parameterization $\vec{\Phi}$.

Def. A **tangent vector** to a manifold, M , at a point $p \in M$, is the tangent vector at p of a curve in M passing through p .



Ex. Find a description of the tangent plane to the torus in \mathbb{R}^4 given by:

$$\vec{\Phi}(u, v) = (\cos u, \sin u, \cos v, \sin v)$$

at the point where: $(u, v) = \left(\frac{\pi}{6}, \frac{\pi}{4}\right)$.

$$D\vec{\Phi}(u, v) = \begin{pmatrix} -\sin u & 0 \\ \cos u & 0 \\ 0 & -\sin v \\ 0 & \cos v \end{pmatrix}$$

$$D\vec{\Phi}\left(\frac{\pi}{6}, \frac{\pi}{4}\right) = \begin{pmatrix} -\frac{1}{2} & 0 \\ \frac{\sqrt{3}}{2} & 0 \\ 0 & -\frac{\sqrt{2}}{2} \\ 0 & \frac{\sqrt{2}}{2} \end{pmatrix}.$$

The tangent space is spanned by the image of $\langle 1, 0 \rangle$ and $\langle 0, 1 \rangle$ under $D\vec{\Phi}\left(\frac{\pi}{6}, \frac{\pi}{4}\right)$.

$$\left(D\vec{\Phi}\left(\frac{\pi}{6}, \frac{\pi}{4}\right)\right) \begin{pmatrix} 1 \\ 0 \end{pmatrix} = \begin{pmatrix} -\frac{1}{2} & 0 \\ \frac{\sqrt{3}}{2} & 0 \\ 0 & -\frac{\sqrt{2}}{2} \\ 0 & \frac{\sqrt{2}}{2} \end{pmatrix} \begin{pmatrix} 1 \\ 0 \end{pmatrix} = \begin{pmatrix} -\frac{1}{2} \\ \frac{\sqrt{3}}{2} \\ 0 \\ 0 \end{pmatrix}$$

$$\left(D\vec{\Phi}\left(\frac{\pi}{6}, \frac{\pi}{4}\right)\right) \begin{pmatrix} 0 \\ 1 \end{pmatrix} = \begin{pmatrix} -\frac{1}{2} & 0 \\ \frac{\sqrt{3}}{2} & 0 \\ 0 & -\frac{\sqrt{2}}{2} \\ 0 & \frac{\sqrt{2}}{2} \end{pmatrix} \begin{pmatrix} 0 \\ 1 \end{pmatrix} = \begin{pmatrix} 0 \\ 0 \\ -\frac{\sqrt{2}}{2} \\ \frac{\sqrt{2}}{2} \end{pmatrix}.$$

We want the tangent space at $\vec{\Phi}\left(\frac{\pi}{6}, \frac{\pi}{4}\right) = \left(\frac{\sqrt{3}}{2}, \frac{1}{2}, \frac{\sqrt{2}}{2}, \frac{\sqrt{2}}{2}\right)$.

So we want all points of the form:

$$\vec{P}(s, t) = \left(\frac{\sqrt{3}}{2}, \frac{1}{2}, \frac{\sqrt{2}}{2}, \frac{\sqrt{2}}{2}\right) + \left(-\frac{1}{2}, \frac{\sqrt{3}}{2}, 0, 0\right)s + \left(0, 0, \frac{\sqrt{2}}{2}, \frac{\sqrt{2}}{2}\right)t$$

where $s, t \in \mathbb{R}$.

Suppose $\vec{\Phi}: U \subseteq \mathbb{R}^k \rightarrow M \subseteq \mathbb{R}^n$ is a parametrization of a k -dimensional manifold M . We know the standard basis for $T_p(\mathbb{R}^k)$, $p \in U$, is given by $(\vec{e}_1)_p, \dots, (\vec{e}_k)_p$. What is the corresponding "standard" basis for $T_{\vec{\Phi}(p)}(M)$?

Ex. Suppose $\vec{\Phi}: U \subseteq \mathbb{R}^2 \rightarrow M \subseteq \mathbb{R}^3$ is a parametrization of a regular surface $M \subseteq \mathbb{R}^3$ given by:

$$\vec{\Phi}(x^1, x^2) = (x(x^1, x^2), y(x^1, x^2), z(x^1, x^2)).$$

By definition, the tangent space, $T_{\vec{\Phi}(p)}(M)$, is the image of:

$$D\vec{\Phi}(p): T_p(\mathbb{R}^2) \rightarrow T_{\vec{\Phi}(p)}(\mathbb{R}^3).$$

So to find the standard basis vectors for $T_{\vec{\Phi}(p)}(M)$, we need to find

$$D\vec{\Phi}(p)(\vec{e}_1)_p \text{ and } D\vec{\Phi}(p)(\vec{e}_2)_p.$$

To calculate this we just need to find the Jacobian matrix for $\vec{\Phi}(p)$:

$$D\vec{\Phi}(p) = \begin{pmatrix} \frac{\partial x}{\partial x^1} & \frac{\partial x}{\partial x^2} \\ \frac{\partial y}{\partial x^1} & \frac{\partial y}{\partial x^2} \\ \frac{\partial z}{\partial x^1} & \frac{\partial z}{\partial x^2} \end{pmatrix}.$$

Now we have:

$$D\vec{\Phi}(p)(\vec{e}_1)_p = \begin{pmatrix} \frac{\partial x}{\partial x^1} & \frac{\partial x}{\partial x^2} \\ \frac{\partial y}{\partial x^1} & \frac{\partial y}{\partial x^2} \\ \frac{\partial z}{\partial x^1} & \frac{\partial z}{\partial x^2} \end{pmatrix} \begin{pmatrix} 1 \\ 0 \end{pmatrix} = \begin{pmatrix} \frac{\partial x}{\partial x^1} \\ \frac{\partial y}{\partial x^1} \\ \frac{\partial z}{\partial x^1} \end{pmatrix} = \frac{\partial \vec{\Phi}}{\partial x^1} = \vec{\Phi}_{x^1}$$

$$D\vec{\Phi}(p)(\vec{e}_2)_p = \begin{pmatrix} \frac{\partial x}{\partial x^1} & \frac{\partial x}{\partial x^2} \\ \frac{\partial y}{\partial x^1} & \frac{\partial y}{\partial x^2} \\ \frac{\partial z}{\partial x^1} & \frac{\partial z}{\partial x^2} \end{pmatrix} \begin{pmatrix} 0 \\ 1 \end{pmatrix} = \begin{pmatrix} \frac{\partial x}{\partial x^2} \\ \frac{\partial y}{\partial x^2} \\ \frac{\partial z}{\partial x^2} \end{pmatrix} = \frac{\partial \vec{\Phi}}{\partial x^2} = \vec{\Phi}_{x^2}.$$

So the standard basis vectors for $T_{\vec{\Phi}(p)}(M)$ are given by $\frac{\partial \vec{\Phi}}{\partial x^1}$ and $\frac{\partial \vec{\Phi}}{\partial x^2}$ evaluated at p .

In general, for a k -dimensional manifold M in \mathbb{R}^n , parametrized by $\vec{\Phi}: U \subseteq \mathbb{R}^k \rightarrow M \subseteq \mathbb{R}^n$, the standard basis for $T_{\vec{\Phi}(p)}(M)$ is $\frac{\partial \vec{\Phi}}{\partial x^1}, \dots, \frac{\partial \vec{\Phi}}{\partial x^k}$ evaluated at p .

Def. Let $U \subseteq \mathbb{R}^k$ be an open set with $p \in U$. Let $\vec{v} \in T_p U$ and $f: U \rightarrow \mathbb{R}$ any differentiable function. We can define a map from the set of real valued, continuously differentiable functions on U , $C^1(U, \mathbb{R})$, into the real numbers by taking a “kind” of **directional derivative** of f at $p \in U$ in the direction of \vec{v}_p by:

$$\vec{v}_p(f) = \left\langle \frac{\partial f}{\partial x^1}, \dots, \frac{\partial f}{\partial x^k} \right\rangle \cdot \left\langle v_1, \dots, v_k \right\rangle$$

where $\frac{\partial f}{\partial x^i}$ is evaluated at p and $\vec{v}_p = v_1(\vec{e}_1)_p + \dots + v_k(\vec{e}_k)_p$. This differs from the standard directional derivative because \vec{v}_p is not necessarily a unit vector.

In particular if $\vec{v}_p = (\vec{e}_i)_p$, then:

$$(\vec{e}_i)_p(f) = \left\langle \frac{\partial f}{\partial x^1}, \dots, \frac{\partial f}{\partial x^k} \right\rangle \cdot \langle 0, \dots, 1, \dots, 0 \rangle = \frac{\partial f}{\partial x^i} \Big|_p.$$

Thus we express $(\vec{e}_1)_p, \dots, (\vec{e}_k)_p$ as: $\frac{\partial}{\partial x^1} \Big|_p, \dots, \frac{\partial}{\partial x^k} \Big|_p$.

Ex. Let $f(x^1, x^2, x^3) = (x^1)^2 + (x^2)(x^3)$.
 Let $p = (-2, 3, 4)$ and $\vec{v}_p = \langle -3, 2, 1 \rangle$.
 Find $\vec{v}_p(f)$.

$$\frac{\partial f}{\partial x^1} = 2x^1 \quad \frac{\partial f}{\partial x^1} \Big|_p = -4$$

$$\frac{\partial f}{\partial x^2} = x^3 \quad \frac{\partial f}{\partial x^2} \Big|_p = 4$$

$$\frac{\partial f}{\partial x^3} = x^2 \quad \frac{\partial f}{\partial x^3} \Big|_p = 3$$

$$(\vec{v}_p)(f) = \langle -4, 4, 3 \rangle \cdot \langle -3, 2, 1 \rangle = 12 + 8 + 3 = 23.$$

We can do the same calculation for manifolds.

If $\vec{\Phi}: U \subseteq \mathbb{R}^k \rightarrow M \subseteq \mathbb{R}^n$, $p \in M$, $\vec{v}_p \in T_p M$, and $f: M \rightarrow \mathbb{R}$, then

$$\vec{v}_p(f) = \left\langle \frac{\partial f}{\partial u^1}, \dots, \frac{\partial f}{\partial u^k} \right\rangle \cdot \langle v_1, \dots, v_k \rangle$$

where (u^1, \dots, u^k) are coordinates on M and:

$$\vec{v}_p = v_1 \vec{\Phi}_{u^1} + \cdots + v_k \vec{\Phi}_{u^k}.$$

Ex. Let $\vec{\Phi}(u^1, u^2) = ((u^1), (u^2), (u^1)^2 + (u^2)^2)$.

Let $(u^1, u^2) = (1, 2)$ so $p = \vec{\Phi}(1, 2) = (1, 2, 5)$.

Let $\vec{v}_p = 3\vec{\Phi}_{u^1}(1, 2) - 2\vec{\Phi}_{u^2}(1, 2)$.

Let $f(x, y, z) = x^4 + y^3 + z^2$. Find $\vec{v}_p(f)$.

We can do this calculation in u^1, u^2 or in x, y, z .

$$\text{In } u^1, u^2: \quad (\vec{v}_p)(f) = \left\langle \frac{\partial f}{\partial u^1}, \frac{\partial f}{\partial u^2} \right\rangle \cdot \langle 3, -2 \rangle$$

$$x = u^1 \quad y = u^2 \quad z = (u^1)^2 + (u^2)^2$$

$$\frac{\partial x}{\partial u^1} = 1 \quad \frac{\partial y}{\partial u^1} = 0 \quad \frac{\partial z}{\partial u^1} = 2u^1$$

$$\frac{\partial x}{\partial u^2} = 0 \quad \frac{\partial y}{\partial u^2} = 1 \quad \frac{\partial z}{\partial u^2} = 2u^2.$$

At p : $x = 1, y = 2, z = 5, u^1 = 1, u^2 = 2$, so we can write:

$$\frac{\partial f}{\partial u^1} = \frac{\partial f}{\partial x} \frac{\partial x}{\partial u^1} + \frac{\partial f}{\partial y} \frac{\partial y}{\partial u^1} + \frac{\partial f}{\partial z} \frac{\partial z}{\partial u^1} = 4x^3(1) + 2z(2u^1)$$

$$\frac{\partial f}{\partial u^1} = 4 + 2(5)(2)(1) = 24$$

$$\frac{\partial f}{\partial u^2} = \frac{\partial f}{\partial x} \frac{\partial x}{\partial u^2} + \frac{\partial f}{\partial y} \frac{\partial y}{\partial u^2} + \frac{\partial f}{\partial z} \frac{\partial z}{\partial u^2} = 3y^2(1) + 2z(2u^2) = 52.$$

$$(\vec{v}_p)(f) = \langle 24, 52 \rangle \cdot \langle 3, -2 \rangle = 72 - 104 = -32.$$

In x, y, z :

$$\begin{aligned}\vec{v}_p &= 3\vec{\Phi}_{u^1}(1, 2) - 2\vec{\Phi}_{u^2}(1, 2) = 3 \langle 1, 0, 2 \rangle - 2 \langle 0, 1, 4 \rangle \\ &= \langle 3, -2, -2 \rangle.\end{aligned}$$

$$\vec{v}_p(f) = \left\langle \frac{\partial f}{\partial x}, \frac{\partial f}{\partial y}, \frac{\partial f}{\partial z} \right\rangle \cdot \langle 3, -2, -2 \rangle$$

$$\frac{\partial f}{\partial x} = 4x^3 \qquad \left. \frac{\partial f}{\partial x} \right|_p = 4$$

$$\frac{\partial f}{\partial y} = 3y^2 \qquad \left. \frac{\partial f}{\partial y} \right|_p = 12$$

$$\frac{\partial f}{\partial z} = 2z \qquad \left. \frac{\partial f}{\partial z} \right|_p = 10$$

$$\vec{v}_p(f) = \langle 4, 12, 10 \rangle \cdot \langle 3, -2, -2 \rangle = 12 - 24 - 20 = -32.$$

Def. A **vector field** on a manifold, M , assigns to each point $x \in M$ a vector in $T_x M$.

Thus, we could let a vector field, \vec{v}_x , on M , map a real valued function, f , on M into a real valued function, g , on M , by $g(x) = \vec{v}_x(f)$.